

学 位 論 文 の 要 旨

氏名 神田翔磨

学 位 論 文 名 Dietary Fiber Deficiency Accelerates Colitis in Mice in the Short Term Independent of Short-Chain Fatty Acids

発 表 雑 誌 名 *Gastroenterology Insights* **15(3)**, 730–743, 2024.
(巻, 初頁～終頁, 年)

著 者 名 Shoma Kanda, Haruki Usuda, Sonoko Karino, Takayuki Okamoto, Tomomi Niibayashi, Takahisa Yano, Kohji Naora, Koichiro Wada

論 文 内 容 の 要 旨

INTRODUCTION

Dietary fiber is considered to keep microbiota healthy state, that may contribute the intestinal barrier function and may increase resistance to colitis-inducing stimuli. In addition, short-chain fatty acids (SCFAs), the bacterial metabolites of dietary fiber, provide intestinal epithelial cells with energy and contribute to maintain epithelial barrier function. Based on these protective property of dietary fiber, intake of dietary fiber is recommended to keep remission of ulcerative colitis. However, protective effects of dietary fiber on colitis are controversial in animals and human. In our present study, we examined whether colitis is caused or worsened under fiber-deficient condition in mice to verify the protective effects of dietary fiber on colitis. In addition, we also examined the protective effect of SCFAs on aggravation of colitis due to fiber deficiency. Furthermore, we evaluated the changes in microbiota due to fiber deficiency to evaluate dysbiosis, that could cause vulnerability in barrier function.

MATERIALS AND METHODS

Mice were administered 2% dextran sulfate sodium (DSS), a colitis inducer, through drinking water. Along with the DSS-administration, mice were fed normal diet or fiber deficient diet. The impact of fiber deficient diet alone without DSS was also examined. Therefore, we made four groups: normal diet without DSS (ND group); fiber-deficient diet without DSS (FF

group); normal diet with DSS (NDD group); fiber deficient diet with DSS (FFD group). To evaluate the severity of colitis, diarrhea, blood in stool, and weight loss were evaluated as macroscopic changes that are associated with colitis. In addition, myeloperoxidase (MPO) activity and histopathological changes in the large intestine were assessed as direct indicators of colitis. To evaluate the involvement of SCFAs in the aggravation of colitis, cecum weight and the amount of SCFAs in the cecum, where SCFAs are produced, were measured. In addition, mixture of SCFAs (the mixture of acetate, propionate and butyrate) were administered orally during the intervention of fiber-deficient diet and DSS, then the colitis-associated parameters were evaluated. Furthermore, the percentages of major bacteria, which comprise more than 1% of the microbiota, were calculated to evaluate the dysbiosis. All experiments with animals in this study were approved by the Animal Care and Use Committee of Shimane University.

RESULTS AND DISCUSSION

The diet feeding and DSS administration were carried out for 5 days (day 0 to day 5). The episodes of diarrhea occurred from day 1 and continued thereafter in the FFD group. There was no diarrhea in NDD group. Blood stool emerged from day 1 and got severer thereafter in the FFD group whereas no or mild blood stool was observed in NDD group. In addition, marked histopathological changes including disruption of epithelium and ulcers were observed in FFD group. In contrast, damage to epithelium and entire villi were mild in NDD group. These abnormal stool state and histopathological changes were not observed both in the ND group and FF group. Furthermore, FFD group showed higher MPO activity compared to NDD group although the MPO activities of both groups were higher than those of ND group and FF group. The MPO activity was not increased in FF group compared to ND group. These results indicate that a fiber deficiency accelerates DSS-induced colitis.

Given the early onset of diarrhea and blood stool in the FFD group, we evaluated the colitis-associated symptoms and direct intestinal damage at day 1 and day 2 to estimate the onset of the aggravation of the colitis. In the comparison of NDD group and FFD group, FFD group showed that enhancement of epithelium damage and MPO activity occurred from day 1 and day 2 respectively. These results suggest that fiber deficiency induced the vulnerability to colitis-inducing stimuli in a very short term.

The amounts of SCFAs in FF group and FFD group were smaller than those in ND group and NDD group. In addition, the amounts of SCFAs between FF group and FFD group, and ND group and NDD group were comparable respectively. The same tendency was observed in the cecum weight. These results indicate that fiber deficiency reduces the amount of SCFAs independent of colitis-inducing stimuli. Nevertheless, the administration of SCFAs did not ameliorate the worsening of the colitis. Therefore, the aggravation of the colitis due to fiber

deficiency would be independent of the decreased amounts of intestinal SCFAs.

In the comparison of the microbiota between the ND group and the FF group, the populations of almost all bacteria were different. The similar tendencies were observed between NDD group and FFD group. These changes were observed day 1 and continued thereafter. Among them, the increased population of *Desulfovibrio* spp. in FF group and FFD group was noteworthy since *Desulfovibrio* spp. are sulfate-reducing bacteria and produce hydrogen sulfide, which likely induce the destruction of epithelial cells as well as upregulation of proteolysis-inducing genes in colonocytes at higher concentration. In addition, the vesicles of the outer membrane destroy the epithelial barrier. Furthermore, *Desulfovibrio* spp. is abundant in IBD patients and positively correlates with the disease activity. Conversely, there have been multiple reports indicating a decrease in the population of *Desulfovibrio* spp. alongside the resolution of colitis in animal models. Therefore, increased *Desulfovibrio* spp. is potential causative factor, which induces vulnerability to DSS-stimulation. Besides, the populations of *Lactobacillus* spp. were lower in FF group and FFD group. Since multiple *Lactobacillus* species have been found to ameliorate DSS-induced colitis, decreased *Lactobacillus* spp. and increase in *Desulfovibrio* spp. instead may have accelerated the formation of unhealthy microbiota that makes intestinal tract irritable to colitis-inducing stimuli.

CONCLUSION

Deficiency of dietary fiber aggravates DSS-induced colitis independent of SCFAs in the short term and the dysbiosis is potential causative factor. These results suggest that a regular intake of dietary fiber is strongly recommended to avoid colitis and preserve intestinal health.